

[AM-09-105] Monte Carlo Simulation

Abstract

Monte Carlo Simulation (MCS) is a computational technique that applies random sampling to model complex systems under uncertainty. In Geographic Information Science and Technology (GIS&T), MCS enables probabilistic analysis of spatial phenomena affected by incomplete data, stochastic processes, or measurement error. Unlike traditional deterministic models that obscure uncertainty, MCS explicitly propagates input variability through simulation, offering robust statistical insights into spatial outcomes. Grounded in statistical principles, MCS supports a wide range of geospatial applications, including flood risk mapping, land use change modeling, satellite image classification, and more. Practical implementation typically involves defining models, characterizing uncertainty through probability distributions, executing simulations, and analyzing the resulting distributions of outcomes. Though there are some limitations, the flexibility and compatibility of modern computing environments have made MCS increasingly accessible, making it a foundational tool for addressing spatial uncertainty and guiding evidence-based policy and analysis.

Keywords: probabalistic, random sampling, simulation, Uncertainty

Author & citation

Oshan, T. M. (2025). Monte Carlo Simulation. The Geographic Information Science & Technology Body of Knowledge (2025 Version), John P. Wilson (Ed.). DOI: [10.22224/gistbok/2025.1.26](https://doi.org/10.22224/gistbok/2025.1.26).

Explanation

1. Introduction
2. Historical Development and Background
3. Theoretical Foundations
4. Practical Implementation
5. Applications in Geospatial Subdisciplines
6. Benefits and Limitations
7. Conclusions

1. Introduction

Monte Carlo Simulation (MCS) is a computational method that uses random sampling to understand and predict the behavior of complex systems, particularly when uncertainty is involved and analytical solutions are impractical. In Geographic Information Science and Technology (GIS&T), MCS has become a vital approach for analyzing spatial phenomena characterized by variability, incomplete data, or stochastic processes (Rubinstein & Kroese, 2016). By simulating numerous realizations using input values drawn from probability distributions, MCS helps quantify uncertainty and generate probabilistic estimates of spatial outcomes

Geospatial data are inherently uncertain due to limitations in measurement precision, spatial resolution, temporal sampling, and modeling assumptions. Traditional deterministic models often mask these uncertainties, providing single-valued outputs that may give a



false impression of precision. In contrast, MCS explicitly incorporates uncertainty by modeling input variables as distributions and examining how these propagate through geospatial analyses (Heuvelink, 1998).

The growing accessibility of high-performance computing, open-source statistical libraries, and spatial modeling tools has further encouraged the adoption of MCS in GIScience applications. These include, but are not limited to, flood risk mapping (Apel et al., 2008), satellite image analysis (Bialek et al., 2020; Wang et al., 2018), urban growth projection (Yeomans & Kozlova, 2023). This entry outlines the theoretical foundations of MCS, describes practical implementation, offers some examples of simulation in geospatial domains, and provides a brief perspective on its potential and limitations.

2. Historical Development and Background

Monte Carlo Simulation was conceived during the 1940s as part of efforts related to nuclear physics and the Manhattan Project. Physicists Stanislaw Ulam and John von Neumann developed the method to estimate complex probabilistic outcomes related to neutron diffusion in fissile materials—problems for which no closed-form solutions were feasible (Metropolis & Ulam, 1949). The method was named after the Monte Carlo Casino in Monaco, as a metaphor for the role of randomness and probability that underpin its design.

The earliest implementations of MCS required manual computation or rudimentary early computers, limiting its initial scope. However, with the advent of modern digital computing in the latter half of the 20th century, Monte Carlo methods gained popularity in fields such as statistical physics, finance, and engineering. By the 1980s and 1990s, improvements in computing power and algorithm design enabled MCS to become a practical tool for many disciplines (Rubinstein & Kroese, 2016), including geospatial analysis.

The integration of MCS into GIS&T was catalyzed by its natural fit for modeling spatial uncertainty. As GIS platforms became programmable and more computationally efficient, researchers began embedding Monte Carlo techniques into spatial decision-support systems, remote sensing workflows, and simulation-based urban planning models (Heuvelink, 1998). The method's capacity to handle nonlinear systems and multivariate uncertainty make it especially appealing for modeling geospatial phenomena.

3. Theoretical Foundations

At the core of Monte Carlo Simulation is the principle of using random numbers to explore complex systems where uncertainty plays a major role. Three key statistical concepts underpin MCS: random sampling, the law of large numbers, and the central limit theorem. Together, these provide a solid mathematical foundation for using MCS in geospatial research and decision-making.

Random sampling is the engine of MCS. Input variables—such as rainfall rates, land surface temperatures, or soil permeability—are represented by probability distributions rather than fixed values. For each simulation run, values are randomly drawn from these distributions. By repeating the model thousands of times with different samples, MCS generates a



distribution of possible outcomes (Kroese et al., 2011). This allows for a rich representation of uncertainty and supports the quantification of risk and variability.

The law of large numbers guarantees that as the number of simulations increases, the average of the results converges to the true expected value. This principle ensures that MCS can provide statistically meaningful outputs, even when input data are uncertain or incomplete (Rubinstein & Kroese, 2016). In practical terms, this means that increasing the number of simulations improves the accuracy of the results, albeit with diminishing returns. The convergence of MCS results can be evaluated through diagnostics such as stability plots or confidence interval narrowing.

The central limit theorem further enhances the utility of MCS by showing that, under general conditions, the distribution of sample means approximates a normal distribution as the number of trials increases, regardless of the underlying input distribution (Zio, 2014). One of the great strengths of Monte Carlo Simulation is its flexibility in accommodating complex, nonlinear relationships among variables. Unlike traditional analytic techniques, MCS imposes no assumptions about linearity or normality in the model structure. This makes it ideal for modeling geospatial systems where interactions are often nonlinear and data may be sparse, noisy, or skewed (Zio, 2014).

In geospatial applications, inputs to Monte Carlo models are often derived from empirical data or expert knowledge and represented as continuous or discrete probability distributions. For example, in flood modeling, rainfall intensity may be modeled using a gamma distribution, while soil moisture parameters could follow a beta distribution. Monte Carlo simulation allows users to define any number of these uncertain inputs, and propagate the variability through a model (Heuvelink, 1998). More advanced simulations may use multivariate distributions to capture correlated uncertainties between spatial variables.

4. Practical Implementation

Monte Carlo Simulation is widely used in GIS&T to support spatial modeling, scenario analysis, and decision-making under uncertainty. The practical implementation of MCS in geospatial workflows typically follows a structured, multi-step process: model formulation, uncertainty characterization, simulation execution, and result analysis.

The first step is model formulation, which involves defining the spatial process or phenomenon to be analyzed. This could range from estimating flood inundation extents to modeling urban land-use change or assessing pollution dispersion. The model should be capable of receiving input parameters as variables rather than fixed values, allowing it to respond dynamically to different scenarios (Heuvelink, 1998).

Next, uncertainty characterization requires identifying the key input variables subject to uncertainty and assigning them appropriate probability distributions. These distributions may be informed by statistical analyses of historical data, published literature, remote sensing observations, or expert knowledge. For example, in an erosion model, rainfall intensity, soil cohesion, and land cover type may all be treated as probabilistic inputs.

In the simulation execution phase, the model is run repeatedly—often thousands of



times—using randomly sampled inputs for each iteration. This generates an ensemble of model outcomes, each reflecting a different realization of the system under study. Software environments such as R and Python (with libraries like NumPy and SciPy), and possibly spatial modeling tools like ArcGIS ModelBuilder or QGIS Processing Toolbox may be used to automate batch simulations.

Once simulations are complete, the final step is analysis of results and interpretation. The outcomes from each iteration are aggregated to form a distribution of results, which is then analyzed statistically. Common outputs include histograms, box plots, quantiles (e.g., 5th and 95th percentiles), and confidence intervals. In geospatial applications, these outputs are often mapped, producing spatial probability surfaces that indicate the likelihood of certain outcomes across a landscape (Heuvelink, 1998).

For example, in flood risk mapping, Monte Carlo simulations can be used to create maps that show the probability of inundation for each pixel or administrative unit. This allows planners to identify high-risk zones and assess the robustness of infrastructure designs under varying hydrological conditions (Apel et al., 2008). In urban growth models, probability maps may indicate which areas are most likely to urbanize under future demographic and policy scenarios (Batty, 2007). Sensitivity analysis is often performed alongside MCS to determine which input variables have the most significant effect on outputs. This information can guide further data collection or model refinement efforts (Kroese et al., 2011).

Practical implementations of MCS in GIS&T also benefit from advancements in cloud computing and parallel processing, which enable larger-scale simulations. Platforms such as Google Earth Engine, Amazon Web Services, and high-performance computing clusters facilitate simulations involving massive spatial datasets and real-time applications.

5. Applications in Geospatial Subdisciplines

5.1 Environmental Modeling

Monte Carlo Simulation is extensively used in environmental modeling to address uncertainty in ecological, hydrological, and atmospheric processes. Environmental systems are complex and driven by numerous interacting variables—many of which are difficult to measure or forecast precisely. MCS offers a way to systematically incorporate this uncertainty and propagate it through environmental models, resulting in more robust predictions and informed decision-making.

In hydrology, MCS is used to assess uncertainties in rainfall inputs, soil infiltration parameters, and watershed responses. For example, researchers may model hundreds or thousands of rainfall-runoff scenarios by sampling from distributions representing rainfall intensity, antecedent moisture conditions, and terrain roughness. The ensemble of results enables planners to understand not only the average response but also the variability of potential flood discharges or groundwater recharge rates (Heuvelink, 1998; Apel et al., 2008).

Ecological modeling can also benefit from MCS, for example, when forecasting the effects of land-use change or climate variability on species distribution. By simulating a range of



possible environmental futures, MCS can estimate the likelihood of habitat suitability or ecosystem collapse, aiding conservation planning.

5.2 Remote Sensing

Remote sensing applications are particularly well-suited to Monte Carlo Simulation due to the inherent uncertainties associated with sensor measurements, atmospheric correction, and image classification. One key application is in the derivation of vegetation indices such as the Normalized Difference Vegetation Index (NDVI). NDVI is computed using reflectance values from the red and near-infrared bands, both of which are subject to calibration error and atmospheric distortion. By applying Monte Carlo techniques—perturbing reflectance values according to estimated sensor noise distributions—researchers can simulate thousands of possible NDVI values for each pixel. The result is a probability distribution rather than a single index value, which can be used to construct uncertainty surfaces across an image (Khalessi et al., 2024).

Land cover classification also benefits from MCS. Classification algorithms often assign each pixel to a land cover category based on decision boundaries that are sensitive to training data and spectral overlap. By simulating multiple realizations of classification based on input variability, researchers can estimate the probability of class membership for each pixel and identify areas of high classification uncertainty (Heuvelink, 1998). These types of uncertain metrics and uncertainty products can enhance decision-making in agriculture, forestry, and environmental monitoring by providing spatially explicit measures of confidence in the remote sensing results.

5.3 Urban Planning

One widely adopted application is urban growth modeling. Cellular automata and agent-based models simulate the expansion of built-up areas over time based on transition probabilities and spatial constraints. MCS can be layered onto these frameworks to reflect uncertainty in transition rules or socio-demographic trends. The resulting ensemble of simulated futures helps identify zones of high agreement where urban expansion is highly probable, and zones of divergence where outcomes are more uncertain (Yeomans & Kozlova, 2023).

Infrastructure planning can also benefit from MCS. For instance, when designing transportation networks, planners must account for future travel demand, fuel prices, and mobility patterns. By using MCS to vary these uncertain inputs, analysts can assess the resilience and efficiency of proposed networks under different assumptions. Monte Carlo-based cost-benefit analyses include the variability of these parameters and others, enhancing the capacity of urban planners to design adaptive, robust strategies that remain effective under a variety of future conditions – an increasingly important goal in a rapidly changing world.

6. Benefits and Limitations

6.1 Benefits

Monte Carlo Simulation offers a number of compelling advantages that have led to its wide



pread adoption across the geospatial sciences. First and foremost is its ability to perform comprehensive uncertainty quantification. By representing uncertain inputs as probability distributions and simulating thousands of model realizations, MCS allows researchers to derive detailed probabilistic insights into spatial phenomena. These outputs include not just mean values, but also variances, percentiles, and full probability distributions—critical for robust decision-making (Rubinstein & Kroese, 2016). A second key benefit is MCS’s model-agnostic flexibility. It can be applied to a wide range of spatial models—from simple overlay analyses in GIS to highly nonlinear hydrological or agent-based simulations. This is particularly important in GIS&T, where diverse data types, modeling techniques, and conceptual frameworks coexist (Kroese et al., 2011). Monte Carlo methods also enable sensitivity analysis, helping analysts determine which inputs contribute most to output variability. This is crucial for prioritizing field measurements, improving model design, and guiding risk communication. For instance, in environmental contamination models, sensitivity analysis data might reveal that uncertainty in soil permeability has a far greater impact on predicted pollutant spread than uncertainty in rainfall inputs. Overall, MCS supports risk-based decision-making, which is increasingly emphasized in policy and planning. Rather than relying on best-case or worst-case scenarios, stakeholders can evaluate the likelihood of outcomes and design strategies that are resilient under a range of plausible futures (Zio, 2014).

6.2 Limitations

Despite its advantages, Monte Carlo Simulation also comes with several limitations that must be acknowledged when applying the technique in GIS&T contexts. The most frequently cited drawback is its computational intensity. Because MCS relies on thousands—or even millions—of model runs to achieve statistical convergence, it can be highly demanding in terms of processing power and time. This is especially challenging when dealing with high-resolution spatial data or computationally expensive models, such as climate or hydrodynamic simulations (Zio, 2014). While parallel processing and cloud computing can mitigate some of these demands, they may not always be accessible. Another important limitation concerns the quality and definition of input distributions. Monte Carlo methods are only as reliable as the assumptions embedded in the input variables. If probability distributions are poorly chosen—due to lack of data, expert bias, or erroneous assumptions—the results can be misleading. In such cases, MCS may give a false sense of precision or understate the true extent of uncertainty (Heuvelink, 1998). Interpretability can also be a challenge. Probabilistic outputs, while richer than single-value estimates, are often harder to communicate to policymakers, stakeholders, and the general public. Concepts like confidence intervals, exceedance probabilities, or distribution tails may require careful explanation and appropriate visual aids to avoid misinterpretation. Finally, MCS is not inherently designed to handle rare but extreme “black swan” events unless these are explicitly built into the model structure or distributions. For example, precipitation values over time may be characterized by a Gaussian distribution, yet the amount of rainfall accumulation during a 100-year storm (i.e., a storm so extreme and rare that it likely does not occur more than once every one hundred years), would be difficult to capture no matter how many simulations were drawn from the distribution that best describes all other precipitation events over a given period. This limits its usefulness in some risk-averse or high-impact scenarios.



7. Conclusion

Monte Carlo Simulation has become a foundational tool in GIS&T for modeling uncertainty, supporting probabilistic decision-making, and enhancing the robustness of spatial analysis. Its theoretical underpinnings in probability theory make it uniquely suited to address the challenges of modeling complex geospatial systems.

Across diverse subfields such as environmental modeling, remote sensing, urban planning, and many more, MCS has proven effective in enhancing deterministic analyses. By producing ensembles of possible outcomes rather than a single prediction, MCS empowers scientists, planners, and decision-makers to assess the range of potential scenarios and make informed choices grounded in statistical evidence.

The increasing availability of high-performance computing, open-source tools, and user-friendly software interfaces has lowered the barrier to entry, making Monte Carlo Simulation more accessible to a wider range of GIS&T professionals. Yet, its implementation must still be approached with care: ensuring accurate input distributions, managing computational demands, and clearly communicating probabilistic results are critical for meaningful application.

As the importance of uncertainty quantification continues to grow in scientific modeling and policy development, the role of Monte Carlo methods in geospatial research will only expand. Future developments are likely to include tighter integration with real-time data, machine learning techniques, and adaptive simulations that adjust based on observed outcomes. In this context, MCS will remain a vital methodology for spatial reasoning in an uncertain world.

References

- [Apel, H., Merz, B., & Thielen, A. H. \(2008\). Quantification of uncertainties in flood risk assessments. *International Journal of River Basin Management*, 6\(2\), 149-162.](#)
- [Batty, M. \(2007\). *Cities and Complexity: Understanding Cities with Cellular Automata, Agent-Based Models, and Fractals*. MIT Press.](#)
- [Białek, A., Vellucci, V., Gentil, B., Antoine, D., Gorroño, J., Fox, N., & Underwood, C. \(2020\). Monte Carlo-Based Quantification of Uncertainties in Determining Ocean Remote Sensing Reflectance from Underwater Fixed-Depth Radiometry Measurements. *Journal of Atmospheric and Oceanic Technology*, 37\(2\), 177-196.](#)
- [Heuvelink, G. B. M. \(1998\). *Error Propagation in Environmental Modelling with GIS*. Wiley](#)
- [Khalesi, F., Ahmed, I., Daponte, P., Picariello, F., De Vito, L., & Tudosa, I. \(2024\). The Uncertainty Assessment by the Monte Carlo Analysis of NDVI Measurements Based on Multi-spectral UAV Imagery. *Sensors*, 24\(9\).](#)
- [Kroese, D. P., Taimre, T., & Botev, Z. I. \(2011\). *Handbook of Monte Carlo Methods*. Wiley.](#)
- [Merz, B., Hall, J., Disse, M., & Schumann, A. \(2010\). Fluvial flood risk management in a changing world. *Natural Hazards and Earth System Sciences*, 10\(3\), 509-527.](#)



[Metropolis, N., & Ulam, S. \(1949\). The Monte Carlo method. *Journal of the American Statistical Association*, 44\(247\), 335-341.](#)

[Rubinstein, R. Y., & Kroese, D. P. \(2016\). *Simulation and the Monte Carlo Method* \(3rd Edition\). Wiley.](#)

[Wang, J., Zhu, J., & Han, X. \(2018\). Using Monte Carlo Simulation to Improve the Performance of Semivariograms for Choosing the Remote Sensing Imagery Resolution for Natural Resource Surveys: Case Study on Three Counties in East, Central, and West China. *ISPRS International Journal of Geo-Information*, 7\(1\).](#)

[Yeomans, J. S., & Kozlova, M. \(2023\). Extending system dynamics modeling using simulation decomposition to improve the urban planning process. *Frontiers in Sustainable Cities*, 5.](#)

[Zio, E. \(2014\). *The Monte Carlo Simulation Method for System Reliability and Risk Analysis*. Springer.](#)

